



TITLE:

<抄録>哺乳動物のフェロモン:雄
えだづのかもしかの耳下腺に含ま
れる活性物質の同定

AUTHOR(S):

桑原, 保正

CITATION:

桑原, 保正. <抄録>哺乳動物のフェロモン:雄えだづのかもしかの耳下
腺に含まれる活性物質の同定. 防虫科学 1974, 39(2): 74-74

ISSUE DATE:

1974-05-31

URL:

<http://hdl.handle.net/2433/158831>

RIGHT:

Summary

Toxic action, acceptability and gustatory effectiveness of the new rodenticide, bisthiosemi [methylene-bis (1-thiosemicarbazide)] with albino rats and Norway rats were investigated.

- (1) The primary action of bisthiosemi may possibly be caused by TSZ being produced by the activation in the stomach.
- (2) It became clear through the preference test, the determination of acceptable thresholds and the acceptance test with the Skinner Box that the acceptability of bisthiosemi in rats was better and larger than that of TSZ.
- (3) By the application of bisthiosemi hypersaturated solution applied on the tongue surface of the rat, no electrical change except the response of off-type was caused in the chorda tympani nerve. Its cause might be related to the extreme insolubility in water.
- (4) The good acceptability of bisthiosemi might be related to both the lack of gustatory effectiveness for the chorda tympani nerve of the

rat and slightly slower toxic action than TSZ.

Acknowledgements: We are indebted to Professor Y. Kawamura and Instructor Y. Kasahara, Department of Oral Physiology, Dental School, Osaka University, for their kindness and guidance, where gustatory effectiveness of bisthiosemi on the chorda tympani nerve of rats was determined. Furthermore, we thank Messrs. A. Hasegawa and Y. Yamamoto at Department of Applied Entomology, Tottori University, for their help in some of the behavior experiments.

References

- 1) Tokumitsu, I., K. Ogushi, H. Yamamoto and T. Kusano: *Botyu-Kagaku*, 38, 202 (1973).
- 2) Kusano, T., Y. Kasahara and Y. Kawamura: *Appl. Ent. Zool.* (in the press).
- 3) Kawamura, Y. and Y. Kasahara: *J. Physiol. Soc. Japan*, 26, 537 (1964) (in Japanese).
- 4) Kusano, T., Y. Kasahara and Y. Kawamura: *Appl. Ent. Zool.*, 6, 40 (1971).
- 5) Nippon Kayaku Co., Ltd.: Technical Information (1972).

抄 録

哺乳動物のフェロモン：雄えだづのかもしかの耳下腺に含まれる活性物質の同定

Mammalian Pheromone: Identification of Active Component in the Subauricular Scent of the Male Pronghorn. Dietland Müller-Schwarze, Christine Müller-Schwarze, Alan G. Singer, Robert M. Silverstein, *Science*, 183, 860~862 (1974).

えだづのかもしかの雄は耳下腺の分泌物をこすり付けて、人間の鼻でもそれと判る匂いを付け、なわ張りのマークとしている。他の雄が付けた匂いを見つけると、あらたにその場所に匂付けを行なう。耳下腺をペンタンで抽出し、揮発性成分をGC分析した結果、8成分が同定された。エステル化後GC-Mass分析により2-methyl butyric acid (1) と isovaleric acid (2) (各々10 μ g/分泌腺) の存在が、13-methyl-1-tetra-

decanol (3) と12-methyl-1-tetradecanol (4) (各々30 μ g/分泌腺) がMass, IR, NMR等の結果同定され、残り4成分(各々70~100 μ g/分泌腺)はメクノリシス後(1)(2)のメチルエステルと(3)(4)を与えること、および機器分析の結果から13-methyl tetradecyl 3-methyl butyrate, 12-methyl tetradecyl 3-methyl butyrate, 13-methyl tetradecyl 2-methyl butyrate, 12-methyl tetradecyl 2-methyl butyrateと同定した。これら8種の化合物およびその混合物の生物試験を行なった結果、isovaleric acidが最も強い活性を示し、その他のエステルやアルコール類は先駆物質、希釈剤あるいは揮散制御剤として働いていると思われる。その他の成分や当該成分の組成化の相異が各個体のマークの識別に役立っていると思われる。

(桑原保正)